



INVEGA TRINZA[®]

paliperidone palmitate extended-release
injectable suspension

273 mg, 410 mg, 546 mg, 819 mg

DISCOVER
INVEGA TRINZA[®]

THE **ONLY** SCHIZOPHRENIA TREATMENT
TO CONTROL SYMPTOMS FOR
THREE MONTHS WITH ONE DOSE.*



*After being adequately treated with INVEGA SUSTENNA[®] (paliperidone palmitate), the once-monthly version of the active medication in INVEGA TRINZA[®], for at least four months.

INDICATION

INVEGA TRINZA[®] (3-month paliperidone palmitate) is a prescription medicine given by injection every 3 months by a healthcare professional and used to treat schizophrenia. INVEGA TRINZA[®] is used in people who have been treated with INVEGA SUSTENNA[®] (1-month paliperidone palmitate) for at least 4 months.

IMPORTANT SAFETY INFORMATION

INVEGA TRINZA[®] can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). INVEGA TRINZA[®] is not approved for treating dementia-related psychosis.

Please see additional Important Safety Information and accompanying full Prescribing Information, including Boxed WARNING.

Image is being used for illustrative purposes only; any person depicted in the image is a model.

ABOUT INVEGA TRINZA[®]

Image is being used for illustrative purposes only; any person depicted in the image is a model.

TAKE CONTROL



It's the only schizophrenia treatment proven to control symptoms for three months with a single dose.

It's used only after you have been adequately treated with INVEGA SUSTENNA[®] (paliperidone palmitate), the once-monthly version of the active medication in INVEGA TRINZA[®], for at least four months.

**ONLY
INVEGA TRINZA[®]**

CONTROLS SYMPTOMS
OVER THREE MONTHS IN
A SINGLE DOSE

&

IS GRADUALLY RELEASED
IN YOUR BODY

&

IS GIVEN ONLY FOUR TIMES
A YEAR

Important Safety Information

Do not receive INVEGA TRINZA[®] if you are allergic to paliperidone palmitate, risperidone, or any of the ingredients in INVEGA TRINZA[®]. See Patient Information leaflet for a complete list of ingredients in INVEGA TRINZA[®].


Please see Important Safety Information enclosed.

TREATMENT IN ACTION



Image is being used for illustrative purposes only; any person depicted in the image is a model.

ONLY FOUR TIMES



Once you start on INVEGA TRINZA®, you only have to get a dose of your medication by injection from a healthcare professional four times a year. INVEGA TRINZA® is gradually released in your body. One dose is proven to control schizophrenia symptoms over three months.

GET RESULTS



In a long-term study, 93% of patients treated with INVEGA TRINZA® did not experience a significant return of schizophrenia symptoms vs. those not treated.

Important Safety Information

Take INVEGA TRINZA® exactly as your healthcare provider tells you to. Your healthcare provider will tell you how much INVEGA TRINZA® you will receive and when you will receive it. INVEGA TRINZA® is a prescription medicine given by injection by a healthcare professional and used to treat schizophrenia.

Do not receive INVEGA TRINZA® if you are allergic to paliperidone palmitate, risperidone, or any of the ingredients in INVEGA TRINZA®. The most common side effects of INVEGA TRINZA® include: injection site reactions, weight gain, headache, upper respiratory tract infections, feeling restlessness or difficulty sitting still, slow movements, tremors, stiffness and shuffling walk. Tell your healthcare provider if you have any side effect that bothers you or does not go away.

GETTING STARTED

INVEGA TRINZA[®] is gradually released in your body. One dose, given by injection from a healthcare professional, is proven to control schizophrenia symptoms for three months. So it only has to be taken four times a year.

If you and your doctor decide that INVEGA TRINZA[®] is right for you, here is how your treatment plan might look.

Important Safety Information

The most common side effects of INVEGA TRINZA[®] include: injection site reactions, weight gain, headache, upper respiratory tract infections, feeling restlessness or difficulty sitting still, slow movements, tremors, stiffness and shuffling walk. Tell your healthcare provider if you have any side effect that bothers you or does not go away.

STARTER DOSES



Month 1

- 1 Your doctor may give you an oral test dose to determine if you are allergic to any of the ingredients of INVEGA TRINZA[®]. Then you will get your first starter dose of once-monthly INVEGA SUSTENNA[®] (paliperidone palmitate), followed by a second starter dose one week later.

Starter doses are not necessary if you are currently taking INVEGA SUSTENNA[®].


ADJUSTING



Month 2 Month 3 Month 4 Month 5

- 2 You will get an injection of INVEGA SUSTENNA[®] for a minimum of four months in a row until your doctor determines the right dose for you.

MAINTENANCE



Month 6 Month 7 Month 8 Month 9 Month 10

- 3 When your doctor tells you it is time, you will be transitioned to the appropriate dose of INVEGA TRINZA[®]. You will then get a dose of INVEGA TRINZA[®] once every three months.

Please see Important Safety Information enclosed.



**MAKE AN
INFORMED DECISION**

Image is being used for illustrative purposes only; any person depicted in the image is a model.

**TALK TO YOUR DOCTOR OR TREATMENT TEAM
IF YOU HAVE ANY QUESTIONS**

**What is the most important information I should
know about INVEGA TRINZA[®]?**

**INVEGA TRINZA[®] can cause serious side effects,
including:**

**Increased risk of death in elderly people who
are confused, have memory loss and have
lost touch with reality (dementia-related
psychosis). INVEGA TRINZA[®] is not for treating
dementia-related psychosis.**

What is INVEGA TRINZA[®]?

INVEGA TRINZA[®] is a prescription medicine given by injection by a healthcare professional and used to treat schizophrenia.

INVEGA TRINZA[®] is used in people who have been treated with INVEGA SUSTENNA[®] 1 time a month injections for at least 4 months.

It is not known if INVEGA TRINZA[®] is safe and effective in children under 18 years of age.

Who should not receive INVEGA TRINZA[®]?

Do not receive INVEGA TRINZA[®] if you:

- are allergic to paliperidone palmitate, risperidone, or any of the ingredients in INVEGA TRINZA[®]. See the end of the Patient Information leaflet for a complete list of ingredients in INVEGA TRINZA[®].

What should I tell my healthcare provider before receiving INVEGA TRINZA®?

Before you receive INVEGA TRINZA®, tell your healthcare provider about all your medical conditions, including if you:

- have had Neuroleptic Malignant Syndrome (NMS)
- have or have had heart problems, including a heart attack, heart failure, abnormal heart rhythm, or long QT syndrome
- have or have had low levels of potassium or magnesium in your blood
- have or have had uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia)
- have or have had kidney or liver problems
- have diabetes or have a family history of diabetes
- have had a low white blood cell count
- have had problems with dizziness or fainting or are being treated for high blood pressure
- have or have had seizures or epilepsy
- have any other medical conditions
- are pregnant or plan to become pregnant. It is not known if INVEGA TRINZA® will harm your unborn baby.
 - If you become pregnant while taking INVEGA TRINZA®, talk to your healthcare provider about registering with the National Pregnancy Registry for Atypical Antipsychotics. You can register by calling 1-866-961-2388 or visit <http://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/>
 - Infants born to women who are treated with INVEGA TRINZA® may have withdrawal symptoms or other symptoms such as tremors, muscle spasms, abnormal movement of arms and legs, and twitching of eyes.

- are breastfeeding or plan to breastfeed. INVEGA TRINZA® can pass into your breast milk and may harm your baby. You and your healthcare provider should decide if you will receive INVEGA TRINZA® or breastfeed. You should not do both.

Patients (particularly the elderly) taking antipsychotics with certain health conditions or those on long-term therapy should be evaluated by their healthcare provider for the potential risk of falls.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Know the medicines you take. Keep a list of them to show to your healthcare provider or pharmacist when you get a new medicine.

How will I receive INVEGA TRINZA®?

- Follow your INVEGA TRINZA® treatment schedule exactly as your healthcare provider tells you to.
- Your healthcare provider will tell you how much INVEGA TRINZA® you will receive and when you will receive it.
- INVEGA TRINZA® is given as an injection by your healthcare provider into the muscle (intramuscularly) of your arm or your buttocks, 1 time every 3 months.

What should I avoid while receiving INVEGA TRINZA®?

- INVEGA TRINZA® may affect your ability to make decisions, think clearly, or react quickly. **Do not** drive, operate heavy machinery, or do other dangerous activities until you know how INVEGA TRINZA® affects you.
- Avoid getting overheated or dehydrated.

What are the possible side effects of INVEGA TRINZA®?

INVEGA TRINZA® may cause serious side effects, including:

- See **“What is the most important information I should know about INVEGA TRINZA®?”**
- **stroke in elderly people (cerebrovascular problems) that can lead to death**
- **Neuroleptic Malignant Syndrome (NMS).**
NMS is a rare but very serious problem that can happen in people who receive INVEGA TRINZA®. NMS can cause death and must be treated in a hospital. Call your healthcare provider right away if you become severely ill and have any of these symptoms:
 - high fever
 - severe muscle stiffness
 - confusion
 - loss of consciousness
 - changes in your breathing, heartbeat and blood pressure
- **problems with your heartbeat.** These heart problems can cause death. Call your healthcare provider right away if you have any of these symptoms:
 - passing out or feeling like you will pass out
 - dizziness
 - feeling as if your heart is pounding or missing beats
- **uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia)**

- **metabolic changes.** Metabolic changes may include high blood sugar (hyperglycemia), diabetes mellitus and changes in the fat levels in your blood (dyslipidemia), and weight gain.
- **low blood pressure and fainting**
- **changes in your blood cell counts**
- **high level of prolactin in your blood (hyperprolactinemia).** INVEGA TRINZA® may cause a rise in the blood levels of a hormone called prolactin (hyperprolactinemia) that may cause side effects including missed menstrual periods, leakage of milk from the breasts, development of breasts in men, or problems with erection.
- **problems thinking clearly and moving your body**
- **seizures**
- **difficulty swallowing that can cause food or liquid to get into your lungs**
- **prolonged or painful erection lasting more than 4 hours.** Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours.
- **problems with control of your body temperature especially when you exercise a lot or spend time doing things that make you warm. It is important for you to drink water to avoid dehydration.**

The most common side effects of INVEGA TRINZA®

include: injection site reactions, weight gain, headache, upper respiratory tract infections, feeling restlessness or difficulty sitting still, slow movements, tremors, stiffness and shuffling walk.

Tell your healthcare provider if you have any side effect that bothers you or does not go away.

These are not all the possible side effects of INVEGA TRINZA®. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

General information about the safe and effective use of INVEGA TRINZA®

Medicines are sometimes prescribed for purposes other than those listed in the Patient Information leaflet. Do not use INVEGA TRINZA® for a condition for which it was not prescribed. Do not give INVEGA TRINZA® to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about INVEGA TRINZA® that is written for health professionals.

The Patient Information leaflet summarizes the most important information about INVEGA TRINZA®. If you would like more information, talk with your healthcare provider.

You can ask your healthcare provider or pharmacist for more information that is written for healthcare professionals. For more information, go to www.invegaTrinza.com or call 1-800-526-7736.

What are the ingredients in INVEGA TRINZA®?

Active ingredient: paliperidone palmitate

Inactive ingredients: polysorbate 20, polyethylene glycol 4000, citric acid monohydrate, sodium dihydrogen phosphate monohydrate, sodium hydroxide, and water for injection

Manufactured by: Janssen Pharmaceuticals, Inc.
Titusville, NJ 08560

© Janssen Pharmaceuticals, Inc. 2017

045409-170321



INVEGA TRINZA[®]

paliperidone palmitate extended-release
injectable suspension

273 mg, 410 mg, 546 mg, 819 mg

**INVEGA TRINZA[®]
IS A CHOICE
MADE TOGETHER**

Image is being used for illustrative purposes only; any person depicted in the image is a model.

FOCUS ON YOUR FUTURE

It's important to keep in mind that not all medications are the same and their effects are different for everybody. Talk to your doctor to decide together which medication is right for you.

Discuss whether you're interested in trying something new and if you have any concerns with your current medication. Everyone's needs are different. Talk to your doctor to see if INVEGA TRINZA[®] fits yours.

Important Safety Information

The most common side effects of INVEGA TRINZA[®] include: injection site reactions, weight gain, headache, upper respiratory tract infections, feeling restlessness or difficulty sitting still, slow movements, tremors, stiffness and shuffling walk.

QUESTIONS TO ASK YOUR DOCTOR

How is INVEGA TRINZA[®] different from my current medication?

How can a single dose work to treat my symptoms over three months?

With my symptoms under control, what goals can I set for my treatment?

Why do I need to take INVEGA SUSTENNA[®] (paliperidone palmitate) before taking INVEGA TRINZA[®]?

What are the possible side effects?

What if I miss a dose?

MAKING PRESCRIPTIONS AFFORDABLE

The INVEGA TRINZA[®] Instant Savings Program is an insurance co-pay support program that helps reduce the patient out-of-pocket cost of INVEGA TRINZA[®] prescriptions. Private or commercially insured patients who qualify will:

- Pay no more than \$10 for each prescription of INVEGA TRINZA[®] purchased through a pharmacy
- Pay no more than \$50 per prescription for medication purchased at the physician's office

Program exhausts after 4 doses of INVEGA TRINZA[®] or \$6000, whichever comes first.

Card not available to individuals enrolled in federally subsidized healthcare programs that cover prescription drugs, including Medicare, Medicaid, TRICARE, or any similar federal or state programs, including patient assistance programs.

This offer may not be combined with any other coupon, discount, prescription savings card, free trial, or other offer.

Participants certify that they will not seek reimbursement or compensation from any of these programs, to include a flexible spending account, a Health Savings Account (HSA), or a Health Reimbursement Account (HRA).

The selling, purchasing, trading, or counterfeiting of this card is prohibited.

Offer good only in the U.S. and Puerto Rico. For Massachusetts residents only, this offer is subject to change per state legislation. Janssen Pharmaceuticals, Inc., reserves the right to rescind, revoke, or amend this offer without notice at any time. Void where prohibited, taxed, or otherwise restricted by law.

Program expires December 31, 2017.

As a condition of participating in this program, you must ensure that you comply with any co-payment disclosure requirements of your insurance carrier or third-party payer, including disclosing to your insurer the amount of co-payment support you receive from this program.

For additional information on INVEGA SUSTENNA[®] go to www.InvegaSustenna.com.


GOVERNMENT INSURANCE OR NO INSURANCE

If you have government insurance or no insurance, visit JanssenPrescriptionAssistance.com for information about other financial assistance.

Find out if you qualify for this insurance co-pay support program by calling 888-535-2850.

Janssen Pharmaceuticals, Inc., reserves the right to rescind, revoke, or amend this offer without notice at any time.

Please see Important Safety Information enclosed.



INDICATION

INVEGA TRINZA® (3-month paliperidone palmitate) is a prescription medicine given by injection every 3 months by a healthcare professional and used to treat schizophrenia. INVEGA TRINZA® is used in people who have been treated with INVEGA SUSTENNA® (1-month paliperidone palmitate) for at least 4 months.

IMPORTANT SAFETY INFORMATION

INVEGA TRINZA® can cause serious side effects, including an increased risk of death in elderly people who are confused, have memory loss, and have lost touch with reality (dementia-related psychosis). INVEGA TRINZA® is not approved for treating dementia-related psychosis.

Please see additional Important Safety Information and accompanying full Prescribing Information, including Boxed WARNING.